

20. Valkova D., J. Encheva, V. Encheva, P. Shindrova, M. Christov (2015). Study of hybrid material originated from interspecific crosses with wild *Helianthus annuus* L. for resistance to diseases and parasite broomrape. Plant Science, vol. LII, No. 4, 18-21.
21. Valkova, D. (2013). Investigation of the species from genus *Helianthus* as sources of important breeding characters. Ph.D. dissertation, 200 p (In Bg).
22. Yasin A. B. and Singh S. (2010). Correlation and path coefficient analyses in sunflower. J. of Plant Breeding and Crop Sci. 2(5): 129-133.

VARIABILITY OF THE MOST IMPORTANT QUANTITATIVE PROPERTIES IN SOME VARIETIES OF TOBACCO TYPE BURLEY

**Milan Mitreski¹, Ana Korubin-Aleksoska¹, Jane Aleksoski², Jordan Trajkoski¹, MarjanTrajkoski³,
Petre Taskoski¹**

¹University “St. Kliment Ohridski” – Bitola, Scientific tobacco Institute-Prilep

²Bis Promet Agrocenter - Bitola

³SEKE-Prilep

Corresponding author: anakorubin@yahoo.com

Abstract

The morphological properties that characterize the type of tobacco or the variety in one type are known as qualitative and quantitative. The quantitative properties are regularly conditioned from the impact of a larger number of genes and also they depend from the environment factors. The investigations are done in 2015 on experimental field in the Scientific Tobacco Institute in Prilep, with four repetitions with following varieties: Posejdon (control), BD-1, B-1246 and one new perspective line DP-1710. The aim of this investigation is to show the variability of the most important quantitative properties: height of the plant with inflorescence, the number of leaves per plant and the length and width of the biggest and the smallest leaf from the middle harvesting belt in tobacco type Burley. The results from the research are processed statistically by these parameters: average value (\bar{x}), average value error ($S\bar{x}$), standard deviation (S), variation coefficient (CV) and variation width (WV). From the research, we established that the subject varieties are stable enough, the variability is very low because everywhere the variation coefficient was lower than 10%. However, the newly created line DP -1710 is with the slightest variation in the tested properties. Among other things, it is on average the highest (\bar{x} = 188 cm) and has the largest leaves (\bar{x} = 63 cm), which is a positive feature in coarse tobacco of this type.

Keywords: tobacco, type Burley, quantitative properties, variability.

Introduction

Tobacco belongs to the group of strategic crops in the agricultural economy in the Republic of Macedonia. The cultivation of tobacco is conducted on an area between 12.000 and 15.000 ha with an annual production of 20 to 25 million kilograms of tobacco raw material with good quality. From the aspect of the typical representation, over 95% from this fields belong to aromatic oriental tobacco type Prilep and Yaka, while the large ones (Virginia and Burley) almost cannot be seen in the fields. By the end of the 90s of the last century these types were grown here which reduced the tobacco raw material import. But they were necessary for the production of the most popular “american blend” cigarettes. In this cigarette, the type Burley is involved with 35% (Mickovski, 2004). According to Devcic et al. (1984), the largest manufacturer in the world of this tobacco type are USA, Mexico, Japan, Italy and Spain. We need to point out that in Macedonia there are areas with perfect conditions for growing tobacco type Burley with good quality, so this makes a challenge for this type to be brought again in the production. Lately, the scientists from the Scientific Tobacco Institute - Prilep are picking and creating types that can meet all needs of the cigarette factories. That’s why the subject and the purpose of this research is the variability of the most quantitative properties of the tobacco type Burley which have good combinational abilities and this is a condition for making new more productive and more qualitative from already existing. This is the only way to bring back the interest for this tobacco type and the factories will have no problem to absorb the tobacco raw

material and plus the financial effects will be guaranteed for everyone in the tobacco industry in the Republic of Macedonia.

Material and methods

The researches were made on three varieties and one new line from the type Burley: Posejdon control variety (\emptyset), BD-1, B-1246 and the line DP-1710. The first three varieties are originally from other countries and some of them are still growing in their homelands and in other countries too (Posejdon was grown in Macedonia before). Despite now they are represented on smaller fields, they are still current because of its good quality and they are also used for selection about this tobacco type. The control type Posejdon (Photo 1) is created from Igor Bolsunov in the research station in Fürstenfeld in Austria. The type BD-1 (Photo 2) originally comes from Serbia and was created from Dobrivoje Jovanovik, the leader of the Department of Genetics and Selection within the Development Sector in Tobacco Industry in Nis. The type B-1246 (Photo 3) comes from Bulgaria and until recently it was fairly represented in the tobacco producing areas because of the recognizable quality of the obtained tobacco raw material. The new line DP-1710 (Photo 4) was created in the Scientific Tobacco Institute - Prilep by crossing and selecting of two Burley tobacco varieties. The experiment was set in the Experimental Field of the Scientific Tobacco Institute - Prilep in 2015, on a deluvial - colluvial soil in four repetitions. During the vegetation, the necessary agrotechnical operations are carried out to ensure normal growth and development of plants (feeding with nitrogen fertilizer, trampling and treating tobacco according to the program of the Tobacco Institute - Prilep for the protection of tobacco from diseases, pests and weeds). The tobacco in the experiment is sprinkled three times with a floating norm of 400 m³ of water per hectare. We need to point out that the production of 2015 was assessed as a good year for tobacco production. Measurement of the quantitative properties (height of the plant with inflorescence, number of leaves and length and width of the largest and smallest leaf of the middle belt of the plant) were carried out in the field in the phase of full blossoming of tobacco according to standard methods in the selection, (the mean value \bar{x}) for each property is determined on the basis of fifteen randomly selected plants of each variety in the experiment. The obtained data from the measurements are statistically processed through parameters of variability of properties (Najcevska, 2002), and the results are shown in tables.



Photo 1. Posejdon



Photo 2. BD-1



Photo 3. B-1246



Photo 4. L. DP-1710

Results and discussion

The investigated quantitative properties of the Burley type tobacco varieties are also known as morphological. They have great meaning in tobacco selection and genetics because they determinate the type and the varieties that belongs to him. Despite this, the number and the size of the leaves are determined the yield and the quality of tobacco. They are dictated by their own

genotype, but are also dependent on soil-climatic conditions in the region where tobacco is grown, as well as from applied agrotechnics during vegetation. We mentioned that the results from the research are shown in table, separately for each property just to be shown better, compared with the examined varieties and the new line and also coming up with a conclusion.

Height of the plant with inflorescence

The height of the tobacco plant is a varieties feature. According this property, Uzunoski (1985), separated tobacco varieties in three groups:

1. Variety with low growth, its height is up to 70 cm (Prilep, low Otlja);
2. Varieties with middle growth, with plant height with the inflorescence from 70 - 130 cm (Yaka, Dzebel);
3. Varieties with high growth, its height is over 130 cm (Virdzinija and Burley). Risteski et al. (2007), made a research on six varieties and they point out that the height of the plant with inflorescence approximately goes from 142 cm for variety B1317 to 194 cm for the variety B-98/N CMS.

The results for this quantitative property are shown in Table 1.

Table 1. Height of the plant with inflorescence (cm)

Varieties	n	\bar{x}	$S\bar{x}$	S	CV (%)	WV
Posejdon \emptyset	15	180	0.89	3.44	1.91	175-187
BD-1	15	172	0.88	3.42	1.99	164-175
B-1246	15	168	0.96	3.70	2.20	163-174
L. DP-1710	15	188	0.63	2.45	1.30	185-195

Number of leaves of the plant

The number of leaves of the plant depends on its genetic structure of the variety and the growing conditions. It is considered that the number of leaves is one of the most stable tobacco quantitative properties. According to Atanasov (1972), the number of leaves is a variety feature and represents highly consistent quantitative property. Dyulgerski (2016), examined the new lines from the type Burley in Bulgaria and confirmed that the highest number of leaves has line 1540 (average 31.6) and the lowest number of leaves has line 1525 (26.3 leaves per stalk). Standard variety Pliska with 27.7 leaves is on the fifth place from all eight examined varieties. Devcic et al. (1984), were researching domestic and foreign tobacco varieties from type Burley in more areas in Croatia and they came to a conclusion that the Posejdon variety has 30 leaves per plant, while Culinec 1 has 28 leaves. Here in our research (Table 2.), e also concluded that Posejdon has the highest number of leaves but we counted 29, DP-1710 has 28, BD-1 and B-1246 are with 27 leaves.

Table 2. Number of leaves of the plant

Varieties	n	\bar{x}	$S\bar{x}$	S	CV (%)	WV
Posejdon \emptyset	15	29	0.29	1.13	3.91	27 - 31
BD-1	15	27	0.30	1.16	4.30	25 - 29
B-1246	15	27	0.34	1.31	4.85	24 - 29
L. DP-1710	15	28	0.20	0.77	2.75	27 - 30

Length of the largest leaf of the plant

The length of the leaves in every tobacco type is very important property because it's in a relationship with the quality of tobacco raw material. In Burley, the larger the leaves are the higher the yield is and the quality is much better. Boceski (2003), points out that the length and the width and also the surface of the leaves during drying is reduced by 20 to 30%, which is very important in the technology of processing the tobacco. Thre results from our measuring and the variability are presented in Table 3.

Table 3. Length of the largest leaf (cm)

Varieties	n	\bar{x}	$S\bar{x}$	S	CV (%)	WV
Posejdon Ø	15	60	0.51	1.96	3.27	56-63
BD-1	15	61	0.47	1.83	3.00	58-63
B-1246	15	59	0.52	2.02	3.41	56-62
L. DP-1710	15	63	0.46	1.77	2.81	60-66

Width of the largest leaf of the plant

The width of the leaves as same as the length depends on the soil-climatic conditions and on the applied agrotechnics during the growing period. Risteski and Kocoska (2013), indicate that the largest leaf of the plant on six examined Burley varieties ranged from 28.2 cm to B-21, up to 40.2 cm in the variety Pelagonec CMS, which had the widest leaves in the experiment. The results that we have from our research are presented in Table 4.

Table 4. Width of the largest leaf (cm)

Varieties	n	\bar{x}	$S\bar{x}$	S	CV (%)	WV
Posejdon Ø	15	30	0.30	1.16	3.87	29-32
BD-1	15	31	0.25	0.96	3.09	30-33
B-1246	15	30	0.31	1.18	3.89	28-32
L. DP-1710	15	32	0.19	0.74	2.27	32-34

Length of the smallest leaf in the middle belt

The leaves of the middle belt of Burley tobacco type are the largest, the highest quality and have a high use value in the processes of processing and fabrication. Therefore, we have measured the length and width of the smallest leaf of this belt in order to determine the number of leaves that can be harvested, taking into account their utilization in the fabrication. In practice it is considered that tobacco dry leaves have length below 25cm and that they are useless, also are increasing production costs, and when purchasing tobacco, they are assessed in the lowest class or are not taken at all by purchasers. Bearing this in mind, it is not recommended harvesting the sum of the top insertions of tobacco of this type. The results of the examination of this property are presented in Table 5.

Table 5. Length of the smallest leaf in the middle belt (cm)

Varieties	n	\bar{x}	$S\bar{x}$	S	CV (%)	WV
Posejdon Ø	15	42	0.37	1.41	3.37	40-45
BD-1	15	43	0.36	1.39	3.32	40-45
B-1246	15	39	0.39	1.51	3.88	36-41
L. DP-1710	15	42	0.27	1.03	2.46	40-44

Width of the smallest leaf of the middle belt

The width of the smallest leaf of the middle belt, along with the length, has an impact on the overall yield and quality of the raw tobacco material from type Burley. The variability of this property is shown by the statistical parameters in Table 6.

Table 6. Width of the smallest leaf of the middle belt (cm)

Varieties	n	\bar{x}	$S\bar{x}$	S	CV (%)	WV
Posejdon Ø	15	21	0.21	0.80	3.79	19-22
BD-1	15	23	0.17	0.64	2.77	22-24
B-1246	15	20	0.22	0.83	4.14	18-21
L. DP-1710	15	22	0.18	0.70	3.19	20-23

References

1. Atanasov, D. (1972). Tobacco Production. ITTI Plovdiv.
2. Boceski, D. (2003). Knowledge and processing of tobacco raw material. Institute for Tobacco - Prilep, II supplemented edition, pp. 210.
3. Devčić, K., Triplat, J., Benković, F. (1984). Sorte Burley Selected at the Tobacco Institute of Zagreb. Tutun/Tobacco, No. 1-2, pp. 3-10.
4. Dyulgovski, Y. (2016). Assessment of biological indicators and production characteristics of perspective lines burley tobacco. Tobacco/Tobacco, 1-6, pp. 13-21.
5. Mickovski, J. (2004). Tobacco in the Republic of Macedonia. Society for Science and Art - Prilep, pp. 162.
6. Najcevska, C. (2002). Experimental statistics applied in agricultural and biological research. Book publishing house Bona-Skopje, pp. 49-71.
7. Risteski, I., Kochoska, K., Hristoski, Zh. (2007). Results of examinations of some foreign and newly created domestic Burley varieties of tobacco in CMS and fertile form. Tobacco / Tobacco, No. 9-10, pp. 200-208.
8. Risteski, I., Kochoska, K. (2013). Results of Comparative Experiment with Berleian Varieties of Tobacco. Report on the Project Tasks of the Scientific Research Work in 2011, University "St. Kliment Ohridski" - Bitola, Scientific Institute for Tobacco - Prilep, pp. 20-24.
9. Uzunovski, M. (1985). Tobacco production. Economic newspaper - Skopje, pp. 96.